

B_s^0 Mixing and Lifetime Difference at CDF

Sinead M. Farrington (for the CDF collaboration)
University of Oxford, Oxford, OX1 3RH, UK

The measurements of the B_s^0 mixing parameter which describes the difference in width (the inverse of the lifetime) between the B_s^0 heavy and light states are described. These measurements include direct measurement of $\Delta\Gamma_s$ and Γ_s by resolving CP eigenstates by angular analysis and measurements in which the proportion of CP even and odd is known.

1. THE B_s^0 MESON SYSTEM

The B_s^0 meson is a bound state consisting of an anti-bottom quark and a strange quark. Its antimatter partner consists of a bottom quark and an anti-strange quark. These two mesons can be interchanged by a second order weak interaction, represented as a Feynman box diagram. The parameters which describe this mixing between B_s^0 and $\bar{B}_s^0$ are predictable parameters in the Standard Model which could be modified by non Standard Model contributions to the box diagram. The system is usually characterised in terms of heavy and light mass eigenstates which are superpositions of the B_s^0 and $\bar{B}_s^0$ and thus there are five key parameters: masses (m_H and m_L), widths (Γ_H and Γ_L) and a phase (β_s). The first measurement of the difference in mass, Δm_s , was made in 2006 [1]. Since this measurement is so precise, the B physics program at CDF now focuses on the measurement of $\Delta\Gamma = \Gamma_L - \Gamma_H$ which is discussed here, β_s and $\Gamma = \Gamma_H + \Gamma_L/2$ discussed in the proceedings of D. Tonelli and S. Behari from this conference.

2. MEASUREMENT OF $\Delta\Gamma$

While Δm_s has been measured to great precision, $\Delta\Gamma_s$ has so far been measured imprecisely. A measurement of $\Delta\Gamma_s$ provides an extra test of the Standard Model since new physics may enter through the phase β_s :

$$\Delta\Gamma = \Delta\Gamma^{SM} \times |\cos 2\beta_s| \quad (1)$$

With measurements of both Δm_s and $\Delta\Gamma$, the following Standard Model relation can be tested, where B , B_s and B_R are parameters from lattice QCD [2]:

$$\frac{\Delta\Gamma_s}{\Delta m_s} = \left[(46.2 \pm 4.4) + (10.6 \pm 1.0) \times \frac{B_s}{B} - (11.9 \pm 1.3) \times \frac{B_R}{B} \right] \times 10^{-4} \quad (2)$$

In interpreting the results presented in this report, the assumption should be made that the B_s^0 light mass eigenstate is CP even and the heavy state is CP odd. Two approaches to measuring $\Delta\Gamma_s$ are pursued. The first is to analyse $B_s^0 \rightarrow J/\psi\phi$ decays, fitting the angular distributions between the decay products in order to decipher the CP odd and even content. The second is to measure the width in a CP specific decay for which the proportion of CP odd and even states is known a priori.

2.1. Direct Measurement of $\Delta\Gamma_s : B_s^0 \rightarrow J/\psi\phi$

Approximately 3150 decays of $B_s^0 \rightarrow J/\psi\phi$ are gathered in 2.8 fb⁻¹ of CDF Run II data. In this sample, a simultaneous fit is made to the mass, lifetime and angular variables. The lifetime distribution and its fit projection are shown in Figure 1. Since the decays are of a pseudo scalar decaying into two vector mesons, the decays contain S and D wave states which are CP even and P wave states which are CP odd. Therefore angular distributions in the J/ψ and ϕ rest frames give information on the CP composition of the $B_s^0 \rightarrow J/\psi\phi$ decays, measured in terms of

$|A_0|$ and $|A_{||}|$ which indicate the S,D and P wave composition. The results of the simultaneous fit assuming no CP violation are

$$\begin{aligned} c\tau_s &= 459 \pm 12(stat) \pm 3(sys) \mu m \\ \Delta\Gamma_s &= 0.02 \pm 0.05(stat) \pm 0.01(sys) ps^{-1} \\ |A_0|^2 &= 0.508 \pm 0.024(stat) \pm 0.008(sys) \\ |A_{||}|^2 &= 0.241 \pm 0.019(stat) \pm 0.007(sys) \end{aligned}$$

The main systematic uncertainties in the $\Delta\Gamma_s$ measurement are from the lifetime resolution model and the background lifetime model. The result can be compared to the predicted value of $\Delta\Gamma_s = 0.096 \pm 0.039 ps^{-1}$ [3]. The previous iteration of this analysis has been published [4].




Figure 1: Lifetime distribution of $B_s^0 \rightarrow J/\psi\phi$ candidates with fit superimposed.

2.2. Measurement of $\Delta\Gamma_s$ from CP specific state: $B_s^0 \rightarrow D_s^+ D_s^-$

The decay $B_s^0 \rightarrow D_s^+ D_s^-$ is a $b \rightarrow c\bar{c}s$ decay which is purely CP even in composition. Therefore a width measurement of $B_s^0 \rightarrow D_s^+ D_s^-$ would measure Γ_L . The branching ratio of the $B_s^0 \rightarrow D_s^+ D_s^-$ mode is related to the difference in width between the two weak eigenstates [5] and so can provide an indirect measurement of $\Delta\Gamma_s$:

$$\frac{\Delta\Gamma_s}{\Gamma_s} = 2 \times BR \left(B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-} \right) \quad (3)$$

The assumption is made that the width difference is mainly caused by $B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-}$ decays. Other decays which are not purely CP even are assumed to have much smaller branching fractions. The $B_s^0 \rightarrow D_s^+ D_s^-$ ($D_s^+ \rightarrow \phi\pi, D_s^- \rightarrow \phi\pi$ or K^*K or $\pi\pi\pi$) branching ratio is measured relative to that of $B_d^0 \rightarrow D_s^+ D^-$ in order to eliminate sources of systematic uncertainty. The remaining candidates with $D_s^+ \rightarrow \phi\pi^+$ after selection are shown in Figure 2 for the $B_s^0 \rightarrow D_s^+ D_s^-$ mode, which is clean owing to the presence of two narrow resonance ϕ mesons in the final state of the D_s^+ decay. The mass distributions are fitted with a three component fit: the Gaussian signal distribution; the combinatorial background distribution obtained by fitting the high sideband and extrapolating it under the peak; and the physics backgrounds, obtained from Monte Carlo simulations. A multi-parameter fit is performed, yielding greater than five standard deviations significance and a branching ratio of [6]:

$$\frac{BR(B_s^0 \rightarrow D_s^+ D_s^-)}{BR(B_d^0 \rightarrow D_s^+ D^-)} = 1.44^{+0.38}_{-0.31}(stat)^{+0.08}_{-0.12}(syst) \pm 0.21 \left(\frac{f_s}{f_d} \right) \pm 0.20(BR(\phi\pi))$$

From this measurement, a 95% confidence level limit of $\Delta\Gamma_s/\Gamma_s \geq 0.012$ can be set using equation 3. A lower limit is set since equation 3 includes excited D_s states while this measurement is made on the non-excited D_s states.

2.3. Measurement of $\Delta\Gamma_s$ from CP specific state: $B_s^0 \rightarrow K^+ K^-$

Fully hadronic decays of B mesons can be resolved at CDF thanks to its displaced track trigger which requires a pair of tracks in the drift chamber at level 1 in the triggering system, and a displaced track in the silicon detector at level 2. In addition, the CDF detector has excellent mass resolution which aids in separation of the various B decays into pairs of hadrons. The projection of the fit to invariant mass, particle identification quantities and a measure of momentum imbalance is shown in Figure 3. The decay $B_s^0 \rightarrow K^+ K^-$ is almost 100% CP even and therefore a




Figure 2: $B_s^0 \rightarrow D_s^+ D_s^-$ candidates with mass fit superimposed.

measurement of the B_s^0 meson lifetime in this state can be combined with knowledge of the average B_s^0 lifetime in order to obtain a measurement of $\Delta\Gamma_s$. A lifetime fit is performed in 0.36 fb^{-1} of CDF Run II data and yields

$$\tau(B_s^0 \rightarrow K^+ K^-) = 1.53 \pm 0.18(\text{stat}) \pm 0.02(\text{sys}) \text{ ps} \quad (4)$$

Using the Heavy Flavour Averaging Group flavour specific lifetime, $\tau(B_s^0) = 1.454 \pm 0.040 \text{ ps}$, yields

$$\Delta\Gamma_s/\Gamma_s = -0.08 \pm 0.23 \pm 0.03 \quad (5)$$




Figure 3: $B_s^0 \rightarrow K^+ K^-$ candidates with mass fit superimposed.

3. CONCLUSIONS

Several measurements of $\Delta\Gamma_s$ have been presented. In the presence of new physics these measurements need not necessarily agree. With increased precision in Tevatron analyses, hints of new physics may be revealed.

References

- [1] A.Abulencia et al., The CDF Collaboration, Phys. Rev. Lett. 97, 242003 (2006)
- [2] A. Lenz, U. Nierste, JHEP 06, 072 (2007)
- [3] A. Lenz, arXiv 0802.0977 (2008)
- [4] T. Aaltonen et al., The CDF Collaboration, Phys. Rev. Lett. 100, 121803 (2008)
- [5] I. Dunietz et al., hep-ph/0012219
- [6] T. Aaltonen et al., The CDF Collaboration, Phys. Rev. Lett. 100, 021803 (2008)